



PMC

US National Library of Medicine
National Institutes of Health[Limits](#) [Advanced](#) [Journal list](#)

Search

[Help](#)[Journal List](#) > [Med Sci Monit](#) > v.19; 2013 > [PMC3718722](#)[PubReader format:
click here to try](#)**Formats:**[Article](#) | [PubReader](#) | [ePub \(beta\)](#) | [PDF \(331K\)](#) | [Citation](#)

Med Sci Monit. 2013; 19: 579–583.

PMCID: PMC3718722

Published online 2013 Jul 15.

doi: [10.12659/MSM.889033](https://doi.org/10.12659/MSM.889033)

Mefloquine use, psychosis, and violence: A retinoid toxicity hypothesis

[Anthony R. Mawson](#)[Author information](#) ▶ [Article notes](#) ▶ [Copyright and License information](#) ▶This article has been [cited by](#) other articles in PMC.**Abstract**Go to:

Mefloquine use has been linked to severe gastrointestinal and neuropsychiatric adverse effects, including cognitive disturbances, anxiety, depression, psychosis, and violence. The adverse effects of the drug are thought to result from the secondary consequences of hepatocellular injury; in fact, mefloquine is known to cause a transient, anicteric chemical hepatitis. However, the mechanism of mefloquine-associated liver damage and the associated neuropsychiatric and behavioral effects of the drug are not well understood. Mefloquine and other 8-amino-quinolines are the only antimalarial drugs that target the liver-stage malaria parasites, which selectively absorb vitamin A from the host. Vitamin A is also stored mainly in the liver, in potentially poisonous concentrations. These observations suggest that both the therapeutic effectiveness of mefloquine and its adverse effects are related to the ability of the 8-aminoquinolines to alter the metabolism of retinoids (vitamin A and its congeners). Several lines of evidence support the hypothesis that mefloquine neurotoxicity and other adverse effects reflect an endogenous form of hypervitaminosis A due to a process involving: mefloquine-induced dehydrogenase inhibition; the accumulation of retinoids in

Related citations in PubMed[Retinoids, race and the pathogenesis of dengue hemorrhagic \[Med Hypotheses. 2013\]](#)[Adverse effects of the antimalaria drug, mefloquine: due \[BMC Public Health. 2002\]](#)[Hypervitaminosis A syndrome: a paradigm of retinoid side \[J Am Acad Dermatol. 1987\]](#)[\[Polymorphism at the MDR1 locus as a cause of mefloquine-induc \[Harefuah. 2010\]](#)[Liver toxicity of retinoid therapy. \[J Am Acad Dermatol. 1988\]](#)[See reviews...](#)[See all...](#)**Cited by other articles in PMC**[Idiosyncratic quinoline central nervous international Journal for Parasitology: Dru...](#)[See all...](#)**Links**[Compound](#)[PubMed](#)**Recent Activity**[Turn Off](#) [Clear](#) [Mefloquine use, psychosis, and violence: A retinoid toxicity hypothesis](#)[See more...](#)

the liver; retinoid-induced hepatocellular damage; the spillage of stored retinoids into the circulation; and the transport of these compounds to the gut and brain in toxic concentrations. The retinoid hypothesis could be tested clinically by comparing cases of mefloquine toxicity and untreated controls in terms of retinoid profiles (retinol, retinyl esters, percent retinyl esters, and retinoic acid). Subject to such tests, retinoid profiling could provide an indicator for assessing mefloquine-associated adverse effects.

Keywords: mefloquine, neurotoxicity, liver, retinoids, depression, psychosis, violence

Background

Go to: 

Mefloquine (Lariam), a synthetic quinoline introduced in the late 1970s for the prevention and treatment of malaria, has been linked to a wide range of adverse cardiovascular, gastrointestinal, and neuropsychiatric effects. Of particular concern are reports of an association between mefloquine use and severe anxiety and depression, cognitive disturbances and, more worryingly, psychosis and violence [1,2]. Mefloquine prophylaxis has been implicated in numerous impulsive suicides and homicides, including deaths in military personnel [3]. After 2002, as a result of such incidents, a history of neuropsychiatric disorder was designated a contraindication for prescribing mefloquine by the U.S. Army. A study carried out in 2007 on a cohort of 11,725 active duty U.S. military personnel showed that 38.4% had received a prescription for mefloquine in the 45 days prior to combat deployment. A review of records showed that 1127 (9.6%) of the entire cohort had contraindications to mefloquine use. Yet 155 persons – 13.8% of those with contraindications and 1.3% of the entire cohort – had been prescribed mefloquine. Thus, 1 in 7 individuals with neuropsychiatric contraindications had received a prescription for mefloquine prior to a recent combat deployment. Since mefloquine continues to be used as an antimalarial, further research is recommended on the incidence and mechanisms of mefloquine-associated adverse events [4].

In a study of the effects of mefloquine on brain function, Hood et al. [5] found that mefloquine caused a concentration-dependent decrease in glutathione and a simultaneous increase in F(2)-isoprostanes in primary rat cortical neurons, indicating the presence of oxidative stress. Mefloquine also induced a concentration-dependent decrease in the number of spines per neuron and spine density, suggesting that oxidative stress caused by the drug was associated with synaptodendritic degeneration.

The molecular mechanisms of mefloquine-induced oxidative stress on brain function and behavior remain uncertain. This paper presents the hypothesis that mefloquine toxicity is an endogenous form of hypervitaminosis A induced by liver damage and the subsequent spillage and transport of stored retinoids in toxic concentration to the brain, gut, and other affected tissues.

Quinolones and Malaria Prophylaxis

Go to:

The quinolones, including chloroquine, primaquine, mefloquine and quinine, are among the most successful yet poorly understood classes of drugs, with important uses in the treatment of systemic lupus erythematosus and HIV as well as malaria. The 8-aminoquinolines in particular (mefloquine, primaquine, and tafenoquine) are the only drugs that target the liver-stage (merozoite) malaria parasites [6]. Malarial infection begins when a person is bitten by an infected female anopheles mosquito and *Plasmodium* parasites are injected into the bloodstream in the form of sporozoites, which travel to the liver. After 7–10 days, during which there are no symptoms, the parasites emerge from the liver cells as merozoites and enter the bloodstream, where they invade and multiply in erythrocytes. When the cells burst, the parasites invade more erythrocytes. Clinical symptoms occur in synchrony with the rupture of infected erythrocytes [7].

Potential clues to understanding both the therapeutic effectiveness of mefloquine and its adverse effects are that the *Plasmodium falciparum* parasite selectively absorbs vitamin A from the host [8] and appears to use the vitamin for its metabolism [9]. Persons with low vitamin A reserves are at increased risk of death from malaria, whereas those with high reserves have less severe disease and are less likely to die from it [10]. Although therapeutically weaker than currently available antimalarial drugs such as artemisinin derivatives, vitamin A has a beneficial adjunctive role in the treatment of malaria and partially protects against malaria infection [11–13]. The selective absorption of vitamin A by the malaria parasite suggests that the effectiveness and toxicity of mefloquine are due in part to the ability of the 8-aminoquinolines to alter the metabolism of vitamin A and its congeners (collectively termed retinoids). As discussed below, there are indications that mefloquine interferes with retinoid metabolism via its effect on dehydrogenase enzymes in the liver.

Retinoids

Go to:

Retinoids are fat-signaling molecules derived mainly from the diet. They are stored principally (about 80% of total vitamin) in the

liver, particularly in the stellate cells, and in sufficient quantity to last the average adult about 2 years without additional intake. In normal physiological concentrations, retinoids are essential for multiple biologic functions, including cellular homeostasis, embryonic development, tissue differentiation and growth, and mucus secretion [14,15]. Retinoic acid (RA), the major active form of vitamin A, binds to and activates specific retinoid receptors that regulate the transcription of many target genes [16]. RA is produced from free retinol in a process that involves the hydrolysis of retinyl esters in the liver, the release of retinol into the circulation, and its subsequent delivery to the target tissues bound to retinol-binding protein (RBP). Retinoic acid is synthesized from the oxidation of retinol to retinaldehyde via an alcohol dehydrogenase, and from retinaldehyde via an aldehyde dehydrogenase reaction [17].

The importance of vitamin A for the nutrition of the parasite suggests that the antimalarial effect of mefloquine could be due to interference with retinoid metabolism by acting as a dehydrogenase inhibitor. This hypothesis is supported by a study in which a functional proteomic approach was used to exploit the structural similarity between quinolones and the purine ring on ATP to identify quinoline-binding proteins. Two human proteins were identified: aldehyde dehydrogenase 1 (ALDH1) and quinone reductase 2 (QR2), and both were found to be selectively inhibited by quinolones. Another inhibitor of ALDH1 (dimethylaminobenzaldehyde) was also lethal to or inhibited the growth of *P. falciparum in vitro* but was less effective in killing the parasites than the quinoline compounds themselves [18]. Chloroquine actively accumulates to millimolar concentrations in the skin and eye when administered at therapeutic levels, and one of the functions of ALDH1 in the eye is to catalyze the conversion of retinaldehyde to retinoic acid [18]. Prolonged use of chloroquine or hydroxychloroquine could therefore cause retinopathy and blindness due to the accumulation of retinaldehyde in the retina [19]. In an earlier related study, mefloquine increased protein degradation but impaired the breakdown of lipids in rat liver lysosomes, resulting in the accumulation of lipids in lysosomes [20].

Mefloquine, Liver Damage, and Retinoids

Go to:

Mefloquine is an effective drug for the prophylaxis and treatment of malaria caused by *P. falciparum*. Although generally well tolerated with few adverse effects, the drug is known to cause a transient, anicteric chemical hepatitis, with slightly elevated liver

enzymes, indicating hepatocellular damage [21]. A review of 516 cases suggested that many of the adverse effects of mefloquine are a post-hepatic syndrome caused by primary liver damage, with symptoms that include malaise, fever, anorexia, headache, abdominal pain, nausea, diarrhea, and concentration difficulties [22]. However, the precise mechanisms linking mefloquine-associated liver damage with neurotoxicity are not well understood.

The hypothesis proposed here is that the action of mefloquine as a dehydrogenase inhibitor leads to the accumulation of retinoids in the liver, resulting in inflammatory changes and subclinical hepatitis. This is followed by the spillage and regurgitation of stored retinoids into the circulation in toxic concentrations and their subsequent transport to the gut and brain, where they induce the adverse gastrointestinal and neuropsychiatric adverse effects of the drug as manifestations of an endogenous form of hypervitaminosis A. The effectiveness of the drug against malaria may be similarly due to hepatic retinoid accumulation and exposure of the merozoite-stage parasites to retinoid concentrations that exceed their tolerance threshold and result in their destruction. Evidence in support for this hypothesis is presented below.

As noted, retinoids are stored in high concentration in the liver, mainly in the form of retinyl esters, and they are normally secreted harmlessly into the circulation as RBP. However, retinyl esters can be extremely toxic if released unbound to protein. An accepted indicator of retinoid toxicity is percent retinyl esters >10% of total vitamin A [23,24]. Liver damage is a known result of excess vitamin A exposure and is associated with reduced serum retinol level but increased liver enzyme levels [25]. Retinoid-induced hepatotoxicity leads to a form of cholestatic liver dysfunction in which bile regurgitates into the circulation, raising the level of all biliary substances in the blood [26,27]. Vitamin A metabolites in bile acids also spill over into the circulation, while stored retinyl esters leak from damaged hepatocytes [28] in proportion to the total liver stores of the vitamin [29]. At the same time, plasma retinol levels decline due to impaired hepatic synthesis and mobilization of the vitamin. The hypothesized net effect is an endogenous form of hypervitaminosis A associated with an increased percentage of plasma retinyl esters as a fraction of total vitamin A, together with increased retinoic acid (RA) concentrations and low or normal concentrations of retinol and its transporter, RBP.

Retinoid Toxicity, Brain and Behavior

Go to: 

The postulated link between mefloquine-induced liver dysfunction and subsequent neuropsychiatric adverse effects due to the spillage of toxic concentrations of retinoids into the circulation receives strong support in the literature. Retinoid signaling pathways affect the functioning of the adult brain via retinal dehydrogenases, cellular retinoid binding proteins, and retinoid receptors, and are essential for cognitive and behavioral development. Furthermore, many neuronal genes are responsive to retinoids, suggesting that retinoid-responsive gene transcription has a significant impact on adult brain function. At normal physiological concentrations in the brain, RA acts as a growth factor, participates in locomotor behavior, and modulates dopaminergic pathways and cells [30]. In the functional cerebral cortex, RA signaling affects the expression and regulation of hundreds of genes through the retinoic acid receptor (RAR) and retinoid X receptor (RXR) classes of ligand-dependent transcription factors, influencing the pattern formation of many organs and tissues [31]. Increased concentrations of RA are found in the adult brain in the striatum and the nucleus accumbens. Retinaldehyde dehydrogenase 1 (RALDH1) is present in the dopaminergic terminals that innervate the striatum from the ventral tegmental area and are necessary for the synthesis of RA in these areas. RA modulates the action of dopamine by regulating the D2 receptor [32].

In higher concentrations, however, retinoids inhibit cell growth and can be pro-oxidant, cytotoxic, mutagenic, and teratogenic. Acute vitamin A toxicity is characterized by nausea, vomiting, headache, vertigo, blurred vision, increased intracranial pressure, irritability, and muscular incoordination. Vitamin A toxicity can occur due to excessive dietary consumption and from treatment with retinoids, as well as endogenously due to medical conditions related to, for example, altered liver and kidney function [24].

Reports have also suggested an association between the use of 13-cis-RA (or isotretinoin), a medication for acne, and the onset of depression, as well as psychosis and suicide in a subgroup of vulnerable individuals [33,34]. This evidence includes case reports, studies showing a temporal association between depression onset and exposure to the drug, challenge and re-challenge cases, evidence of a drug class effect, dose response, and the existence of biologically plausible mechanisms for the association [35–37]. Therapeutic doses reportedly induce cognitive disturbances and depression in from 1% to 11% of patients [38].

Studies in mice also show that 13-cis-RA adversely affects

learning and memory and significantly increases depression-like behaviors [39,40]. Recalling the study of Hood et al. [5], which showed that mefloquine induces oxidative stress on brain function, other reports suggest that acute and chronic vitamin supplementation of laboratory animals at therapeutic doses also leads to increased levels of markers of oxidative stress in liver mitochondria and substantia nigra, and to alterations in locomotor and exploratory activity [41]. Vitamin A supplementation at clinical doses also impairs mitochondrial function in rat hippocampus, decreases brain-derived neurotrophic factor levels and dopamine D2 receptor levels, and decreases glutamate uptake [42].

Similarities Between Vitamin A Poisoning and Impulsive Violence [Go to:](#)

Hypervitaminosis A was unwittingly induced in early Arctic explorers by consuming vitamin A-rich polar bear or seal liver. Reported symptoms included drowsiness, irritability, severe headaches, nausea, and various forms of impulsive and irrational behavior called *pibloktoq* by the Eskimos [43]. Unlike the sympathetic nervous system-based “fight-or-flight” response, in which the senses are heightened, cognitive functions are at their sharpest and the skeletal muscles primed for decisive action, the arousal state associated with hypervitaminosis A is more akin to one of intoxication. Reports of incidents of impulsive violence suggest similarities to those induced by hypervitaminosis A. Such accounts describe violent acts as being carried out in a state of diffuse cognitive impairment, perceptual distortion and anhedonia, and in a dream-like state from which the subject later emerges with little or no memory of the acts in question [44–46].

In addition to causing neuropsychiatric symptoms such as depression, psychosis, and violence, synthetic retinoids in particular have also been linked to a wide range of adverse and often severe gastrointestinal effects [47]. For instance, in a large case-control study involving thousands of patients with Crohn’s disease (CD) and ulcerative colitis (UC), it was found that UC was strongly associated with previous isotretinoin exposure (odds ratio (OR) 4.36, 95% confidence interval (CI): 1.97, 9.66) but not with CD (OR 0.68, 95% CI: 0.28, 1.68). Moreover, increasing the dose of isotretinoin was associated with elevated risk of UC, and the risk of UC was highest in those exposed to isotretinoin for more than 2 months (OR 5.63, 95% CI: 2.10, 15.03) [48]. This evidence suggests that both the adverse neuropsychiatric and gastrointestinal effects of mefloquine may be mediated by retinoid overexpression

and toxicity.

Conclusions

Go to:

The use of mefloquine in the prevention and treatment of malaria has been increasingly linked to a broad range of neuropsychiatric effects, including depression, psychosis, and violence. Mefloquine is known to cause a transient, anicteric chemical hepatitis and the adverse effects of the drug have been postulated to result from the secondary consequences of hepatocellular injury. However, the precise mechanisms linking mefloquine-induced liver damage and the adverse neuropsychiatric effects are not well understood. The explanation for both the therapeutic effectiveness of mefloquine and its adverse consequences may lie in its ability to interfere with the hepatic metabolism of retinol, possibly via dehydrogenase inhibition. This would be expected to cause retinyl esters to accumulate to high concentrations in the liver and also to destroy the merozoite-stage malaria parasites *in situ*. Thus, mefloquine and other 8-amino quinolones may cause liver damage in susceptible persons due to the accumulation of retinoids, which are known to be potentially hepatotoxic. The symptoms of mefloquine toxicity may result from the spillage of stored retinoids from the damaged liver into the circulation and their transport to the gut and brain, causing the adverse neuropsychiatric and gastrointestinal symptoms as a function of an endogenous form of hypervitaminosis A. Indeed, evidence has been presented in this paper that supports this hypothesis. The retinoid hypothesis could be tested clinically by comparing cases of mefloquine toxicity and untreated controls in terms of retinoid profiles (retinol, retinyl esters, percent retinyl esters, and retinoic acid). Cases would be expected to have a significantly increased percentage of plasma retinyl esters as a fraction of total vitamin A, as well as increased retinoic acid (RA) concentrations and low or normal concentrations of retinol and its transporter, RBP. Subject to the outcomes of such tests, retinoid profiling could provide an indicator for assessing mefloquine-associated adverse effects.

Footnotes

Go to:

Source of support: Self financing

References

Go to:

1. Croft AM. A lesson learnt: the rise and fall of Lariam and Halfan. *J Roy Soc Med*. 2007;100:170–74. [[PMC free article](#)] [[PubMed](#)]
2. Toovey S. Mefloquine neurotoxicity: a literature review. *Travel*

Med Infect Dis. 2009;7(1):2–6. [[PubMed](#)]

3. http://www.huffingtonpost.com/2012/03/25/robert-bales-malaria-drug_n_1378671.html?view+politics/

4. Nevin RL. Mefloquine prescriptions in the presence of contraindications: prevalence among US military personnel deployed to Afghanistan, 2007. *Pharmacoepidemiol Drug Saf.* 2010;19(2):206–10. [[PubMed](#)]

5. Hood JE, Jenkins JW, Milatovic D, et al. Mefloquine induces oxidative stress and neurodegeneration in primary rat cortical neurons. *Neurotoxicology.* 2010;31(5):515–23. [[PubMed](#)]

6. Greenwood BM, Fidock DA, Kyle DE, et al. Malaria: progress, perils, and prospects for eradication. *J Clin Invest.* 2008;118(4):1266–76. [[PMC free article](#)] [[PubMed](#)]

7. Clark IA, Cowden WB. The pathophysiology of falciparum malaria. *Pharmacol Ther.* 2003;99:221–60. [[PubMed](#)]

8. Mizuno Y, Kawazu SI, Kano S, et al. *In-vitro* uptake of vitamin A by *Plasmodium falciparum*. *Ann Trop Med Parasitol.* 2003;97(3):237–43. [[PubMed](#)]

9. Guardo I, Lehman LG, Some IT, et al. Influence of malaria on the serum levels of vitamin A, zinc and calcium of children in Doula-Cameroon. *Afr J Biotechnol.* 2007;6:871–876.

10. Friedman JF, Kurtis J, Mtalib R, et al. Malaria is related to decreased nutritional status among male adolescents and adults in the setting of intense perennial transmission. *J Infect Dis.* 2003;188:449–57. [[PubMed](#)]

11. Serghides L, Kain KC. Mechanism of protection induced by vitamin A in falciparum malaria. *Lancet.* 2002;359:1404–6. [[PubMed](#)]

12. Hamzah J, Skinner-Adams TS, Davis TM. *In vitro* antimalarial activity of retinoids and the influence of selective retinoic acid receptor antagonists. *Acta Tropica.* 2003;87:345–53. [[PubMed](#)]

13. SanJoaquin MA, Molyneux M. Malaria and vitamin A deficiency in African children: a vicious circle? *Malaria Journal.* 2009;8:134. [[PMC free article](#)] [[PubMed](#)]

14. Maden M. Retinoid signaling in the development of the central nervous system. *Nature Rev Neurosci.* 2002;3:843–53. [[PubMed](#)]

15. D’Ambrosio DN, Clugston RD, Blaner WS. Vitamin A metabolism: an update. *Nutrients.* 2011;3(1):63–103.

[\[PMC free article\]](#) [\[PubMed\]](#)

16. Theodosiou M, Laudet V, Schubert M. From carrot to clinic: an overview of the retinoic acid signaling pathway. *Cell Mol Life Sci.* 2010;67:1423–45. [\[PubMed\]](#)

17. Blomhoff R, Blomhoff HK. Overview of retinoid metabolism and function. *J Neurobiol.* 2006;66:606–30. [\[PubMed\]](#)

18. Graves PR, Kwiek JJ, Fadden P, et al. Discovery of novel targets of quinoline drugs in the human purine binding proteome. *Mol Pharmacol.* 2002;62:1364–72. [\[PubMed\]](#)

19. Rynes RI. Antimalarial drugs in the treatment of rheumatological diseases. *Br J Rheumatol.* 1997;36(7):799–805. [\[PubMed\]](#)

20. Glaumann H, Motakefi AM, Jansson H. Intracellular distribution and effect of the antimalarial drug mefloquine on lysosomes of rat liver. *Liver.* 1992;12(4 Pt 1):183–90. [\[PubMed\]](#)

21. Gotsman I, Azaz-Livshits T, Fridlender Z, et al. Mefloquine-induced acute hepatitis. *Pharmacotherapy.* 2000;20(12):1517–19. [\[PubMed\]](#)

22. Croft AM, Herxheimer A. Adverse effects of the antimalarial drug, mefloquine: due to primary liver damage with secondary thyroid involvement? *BMC Public Health.* 2002;2:6. [\[PMC free article\]](#) [\[PubMed\]](#)

23. Russell RM. Vitamin and trace mineral deficiency and excess. In: Kasper DL, Braunwald E, Fauci AS, et al., editors. *Harrison's Principles of Internal Medicine.* 16th ed. McGraw-Hill; New York, NY, USA: 2004. pp. 403–11.

24. Penniston KL, Tanumihardjo SA. The acute and chronic toxic effects of vitamin A. *Am J Clin Nutr.* 2006;83:191–201. [\[PubMed\]](#)

25. Alecron-Corredor OM, Alfonso R. Clinical and biochemical alterations in rats treated with high doses of vitamin A. *Arch Latinoam Nutr.* 2007;57(3):224–30. [\[PubMed\]](#)

26. Popper H, Schaffner F. Cholestasis. In: Berk JE, editor. *Gastroenterology.* 4th edit. Vol. 5. W.B. Saunders Co; Philadelphia: 1985. pp. 2697–731. Chapter 148.

27. Ramanathan V, Hensley G, French S, et al. Hypervitaminosis A inducing intra-hepatic cholestasis—a rare case report. *Exp Mol Pathol.* 2010;898:324–25. [\[PubMed\]](#)

28. Leo MA, Lieber CS. New pathway of retinol metabolism in liver microsomes. *J Biol Chem*. 1985;260:5228–31. [[PubMed](#)]
29. Olson JA. Vitamin A – functions, dietary requirements and safety in humans. In: Ziegler EE, Filer LJ Jr, editors. *Present knowledge in nutrition*. 7th ed. Washington, DC: International Life Sciences Institute Press; 2001. pp. 109–19.
30. Lane MA, Bailey SJ. Role of retinoid signaling in the adult brain. *Prog Neurobiol*. 2005;72:275–93. [[PubMed](#)]
31. Luo T, Wagner E, Dräger UC. Integrating retinoic acid signaling with brain function. *Developmental Psychology*. 2009;45(1):139–50. [[PubMed](#)]
32. Krezel W, Ghyselinck N, Samad TA, et al. Impaired locomotion and dopamine signaling in retinoid receptor mutant mice. *Science*. 1998;279:863–67. [[PubMed](#)]
33. Wysowski DK, Pitts M, Beitz J. An analysis of reports of depression and suicide in patients treated with isotretinoin. *J Am Acad Dermatol*. 2001;45(4):515–19. [[PubMed](#)]
34. Hull PR, D’Arcy C. Isotretinoin use and subsequent depression and suicide: presenting the evidence. *Am J Clin Dermatol*. 2004;4:493–505. [[PubMed](#)]
35. Barak Y, Wohl Y, Greenberg Y, et al. Affective psychosis following Accutane (isotretinoin) treatment. *Int Clin Psychopharmacol*. 2005;20:39–41. [[PubMed](#)]
36. Bremner JD, McCaffery P. The neurobiology of retinoic acid in affective disorders. *Prog Neuropsychopharmacol Biol Psychiat*. 2008;32(2):315–31. [[PMC free article](#)] [[PubMed](#)]
37. Kontaxakis VP, Skourides D, Ferentinos P, et al. Isotretinoin and psychopathology: a review. *Ann Gen Psychiat*. 2009;8:2. [[PMC free article](#)] [[PubMed](#)]
38. Bremner JD, Shearer KD, McCaffery PJ. Retinoic acid and affective disorders: the evidence for an association. *J Clin Psychiat*. 2012;73(1):37–50. [[PMC free article](#)] [[PubMed](#)]
39. Crandall J, Sakai Y, Zhang J, Koul O, et al. 13-Cis-retinoic acid suppresses hippocampal cell division and hippocampal-dependent learning in mice. *Proc Nat Acad Sci USA*. 2004;101:5111–16. [[PMC free article](#)] [[PubMed](#)]
40. O’Reilly KC, Shumaker J, Gonzalez-Lima F, et al. Chronic administration of 13-cis- retinoic acid increases depression-related behavior in mice. *Neuropsychopharmacology*. 2006;31:1919–27.

[\[PubMed\]](#)

41. de Oliveira MR, Silvestrin RB, Mello e Souza T, Moreira JC. Therapeutic vitamin A doses increase the levels of markers of oxidative insult in substantia nigra and decrease locomotor and exploratory activity in rats after acute and chronic supplementation. *Neurochem Res.* 2008;33(3):378–83. [\[PubMed\]](#)

42. de Oliveira MR, da Rocha RF, Pasquali MA, Moreira JC. The effects of vitamin A supplementation for 3 months on adult rat nigrostriatal axis: increased monoamine oxidase enzyme activity, mitochondrial redox dysfunction, increased β -amyloid(1–40) peptide and TNF- α contents, and susceptibility of mitochondria to an *in vitro* H₂O₂ challenge. *Brain Res Bull.* 2012;87(4–5):432–44. [\[PubMed\]](#)

43. O'Donnell J. Polar hysteria: an expression of hypervitaminosis A. *Am J Ther.* 2004;11(6):507–16. [\[PubMed\]](#)

44. Bowerman WM. Impaired cognition – a ubiquitous finding in psychiatric disorders. *Orthomol Psychiat.* 1979;8:227–34.

45. Mawson AR. *Transient Criminality: A Model of Stress-Induced Crime.* Praeger Publishers; New York: 1987. pp. 127–54.

46. Mawson AR. Toward a theory of childhood learning disorders, hyperactivity, and aggression. *ISRN Psychiatry.* 2012;2012:589792. [\[PMC free article\]](#) [\[PubMed\]](#)

47. Hanson N, Leachman S. Safety issues in isotretinoin therapy. *Sem Cutan Med Surg.* 2001;20(3):166–83. [\[PubMed\]](#)

48. Crockett SD, Porter CQ, Martin CF, et al. Isotretinoin use and the risk of inflammatory bowel disease: a case-control study. *Am J Gastroenterol.* 2010;105(9):1986–93. [\[PMC free article\]](#) [\[PubMed\]](#)

Articles from *Medical Science Monitor : International Medical Journal of Experimental and Clinical Research* are provided here courtesy of
International Scientific Literature, Inc.

You are here: [NCBI](#) > [Literature](#) > [PubMed Central \(PMC\)](#)

[Write to the Help Desk](#)

GETTING STARTED

[NCBI Education](#)
[NCBI Help Manual](#)
[NCBI Handbook](#)
[Training & Tutorials](#)

RESOURCES

[Chemicals & Bioassays](#)
[Data & Software](#)
[DNA & RNA](#)
[Domains & Structures](#)
[Genes & Expression](#)
[Genetics & Medicine](#)

POPULAR

[PubMed](#)
[Bookshelf](#)
[PubMed Central](#)
[PubMed Health](#)
[BLAST](#)
[Nucleotide](#)

FEATURED

[Genetic Testing Registry](#)
[PubMed Health](#)
[GenBank](#)
[Reference Sequences](#)
[Gene Expression Omnibus](#)
[Map Viewer](#)

NCBI INFORMATION

[About NCBI](#)
[Research at NCBI](#)
[NCBI News](#)
[NCBI FTP Site](#)
[NCBI on Facebook](#)
[NCBI on Twitter](#)

[Genomes & Maps](#)

[Homology](#)

[Literature](#)

[Proteins](#)

[Sequence Analysis](#)

[Taxonomy](#)

[Training & Tutorials](#)

[Variation](#)

[Genome](#)

[SNP](#)

[Gene](#)

[Protein](#)

[PubChem](#)

[Human Genome](#)

[Mouse Genome](#)

[Influenza Virus](#)

[Primer-BLAST](#)

[Sequence Read Archive](#)

[NCBI on YouTube](#)

[Copyright](#) | [Disclaimer](#) | [Privacy](#) | [Browsers](#) | [Accessibility](#) | [Contact](#)

National Center for Biotechnology Information, U.S. National Library of Medicine
8600 Rockville Pike, Bethesda MD, 20894 USA

